arXiv:1610.04979v1 [physics.plasm-ph] 17 Oct 2016

Geometric field theory and weak Euler-Lagrange equation for classical relativistic particle-field systems

Peifeng Fan,[1,2,3] Hong Qin,[2,3,4,*] Jian Liu,[2,5,3] Nong Xiang,[1,3] and Zhi Yu[1,3]

[1]*Institute of Plasma Physics, Chinese Academy of Sciences, Hefei, Anhui 230031, China*

[2]*School of Nuclear Science and Technology and Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China*

[3]*Center for Magnetic Fusion Theory, Chinese Academy of Sciences, Hefei 230031, China*

[4]*Plasma Physics Laboratory, Princeton University, Princeton, NJ 08543, USA*

[5]*Key Laboratory of Geospace Environment, CAS, Hefei, Anhui 230026, China*

Abstract

A manifestly covariant, or geometric, field theory for relativistic classical particle-field system is developed. The connection between space-time symmetry and energy-momentum conservation laws for the system is established geometrically without splitting the space and time coordinates, i.e., space-time is treated as one identity without choosing a coordinate system. To achieve this goal, we need to overcome two difficulties. The first difficulty arises from the fact that particles and field reside on different manifold. As a result, the geometric Lagrangian density of the system is a function of the 4-potential of electromagnetic fields and also a functional of particles' world-lines. The other difficulty associated with the geometric setting is due to the mass-shell condition. The standard Euler-Lagrange (EL) equation for a particle is generalized into the geometric EL equation when the mass-shell condition is imposed. For the particle-field system, the geometric EL equation is further generalized into a weak geometric EL equation for particles. With the EL equation for field and the geometric weak EL equation for particles, symmetries and conservation laws can be established geometrically. A geometric expression for the energy-momentum tensor for particles is derived for the first time, which recovers the non-geometric form in the existing literature for a chosen coordinate system.

PACS numbers: 52.35.Hr, 52.35.-g, 52.35.We, 42.50.Tx, 52.50.Sw

* corresponding author: hongqin@ustc.edu.cn

I. INTRODUCTION

In field theory, it is well-known that energy-momentum conservation laws could be derived from the space-time symmetry of the Lagrangian (or Lagrangian density) that the system admits [1, 2]. For example, the Lagrangian density for the electromagnetic field is

$$\mathcal{L}_F = \frac{1}{8\pi}\left[\left(-\frac{1}{c}\frac{\partial \mathbf{A}}{\partial t} - \nabla\varphi\right)^2 - (\nabla \times \mathbf{A})^2\right]. \tag{1}$$

The energy and momentum conservation laws of the system,

$$\frac{\partial}{\partial t}\left[\frac{\mathbf{E^2} + \mathbf{B^2}}{8\pi}\right] + \nabla \cdot \left[\frac{c}{4\pi}\mathbf{E} \times \mathbf{B}\right] = 0, \tag{2}$$

$$\frac{\partial}{\partial t}\left[\frac{c}{4\pi}\mathbf{E} \times \mathbf{B}\right] + \nabla \cdot \left[\frac{\mathbf{E^2} + \mathbf{B^2}}{8\pi}\mathbf{I} - \frac{\mathbf{EE} + \mathbf{BB}}{4\pi}\right] = 0, \tag{3}$$

can be derived from EL equations,

$$\frac{\partial \mathcal{L}_F}{\partial \varphi} - \frac{D}{D\boldsymbol{x}} \cdot \frac{\partial \mathcal{L}_F}{\partial \dot{\varphi}} = 0, \tag{4}$$

$$\frac{\partial \mathcal{L}_F}{\partial \mathbf{A}} - \frac{D}{D\boldsymbol{x}} \cdot \frac{\partial \mathcal{L}_F}{\partial \nabla \mathbf{A}} - \frac{D}{Dt}\frac{\partial \mathcal{L}_F}{\partial \mathbf{A}_{,t}} = 0, \tag{5}$$

and the symmetry conditions,

$$\frac{\partial \mathcal{L}_F}{\partial t} = 0, \tag{6}$$

$$\frac{\partial \mathcal{L}_F}{\partial \boldsymbol{x}} = 0. \tag{7}$$

Here $D/D\boldsymbol{x}$ and D/Dt denote partial derivative with respect to $\boldsymbol{x}$ and t, respectively, when they operate on a field defined on the space time. Moreover, in relativity, the symmetries of time and space of the Lagrangian density are components of the space-time symmetry,

$$\frac{\partial \mathcal{L}_F}{\partial \chi^\mu} = 0, \quad (\mu = 0, 1, 2, 3), \tag{8}$$

where χ^μ is an arbitrary world point in 4 dimensional Minkowski space-time, i.e., $\chi^0 \equiv t$ and $\chi^i \equiv x^i$. In this paper, we assume that the Minkowski space-time is endowed with a Lorentzian metric and the signature of the metric is $(+---)$ [3]. Equation (8) is a manifestly covariant form of Eqs. (6) and (7). In this paper, the phrase "manifestly covariant" will

be substituted by "geometric", which indicates that covariance is the intrinsic coordinate-independent property of the physical system [4]. Similarly, the EL equations (4) and (5) or the Maxwell equations can be written in the geometric form as

$$\frac{\partial \mathcal{L}_F}{\partial A_\mu} - \frac{D}{D\chi^\nu}\left[\frac{\partial \mathcal{L}_F}{\partial\left(\partial_\nu A_\mu\right)}\right] = 0. \tag{9}$$

Here, the operator $D/D\chi^\nu$ denotes partial derivative with respect to χ^ν while keeping other $\chi^\mu(\mu \neq \nu)$ /fixed, when it is operated on a field on space-time.The geometric energy-momentum conservation law is

$$\partial_\nu T_F^{\mu\nu} = 0, \quad (\mu, \nu = 0, 1, 2, 3) \tag{10}$$

where $T_F^{\mu\nu}$ is the energy-momentum tensor of electromagnetic fields and could be written in a explict form as

$$T_F^{\mu\nu} = \frac{1}{4\pi}\left(-F^{\mu\sigma}F^{\nu}_{\ \sigma} + \frac{1}{4\pi}\eta^{\mu\nu}F_{\sigma\rho}F^{\sigma\rho}\right), \quad (\mu, \nu, \sigma, \rho = 0, 1, 2, 3) \tag{11}$$

where F is electromagnetic tensor and η is the Lorentzian metric. These, of course, are well-known [5, 6].

For classical particle-field systems, where many charged particles evolve under the electromagnetic field generated self-consistently by the particles, the relations between symmetries and conservation laws have only been established recently by Qin et al. [7]. For classical particle-field systems, Qin et al. pointed out that the standard EL equation for particles doesn't hold anymore, because the dynamics of the particles and fields are defined on different manifolds which have different dimensions. The electromagnetic fields are defined on the space-time domain, whereas the particle trajectories as fields are only defined on the time-axis. For particles, a weak EL equation is established to replace the standard EL equation. It was discovered that the weak EL equation can also link symmetries with conservation laws as in standard field theory. This field theory is non-relativistic, but it can be easily extended to relativistic cases, which will be given in Sec. II. However, this approach is based on the split form of space and time. In another word, it is not geometric.

In this paper, we will geometrically reformulate the field theory for classical particle-field system established by Qin et al. [7]. A geometric weak EL equation will be derived, and the energy-momentum conservation will be geometrically derived from the space-time symmetry. To achieve this goal, we need to overcome two difficulties. The first difficulty arises from

the fact that particles and field reside on different manifold as noticed in Ref. [7]. In the geometric setting, this difference is more prominent. Particles' dynamics are characterized by world-lines on the space-time. They are defined on R^1 as a field valued in the space-time, and the domain of the particle fields can be the proper time or any other parameterization for the world-lines. The world-line of a particle is uniquely defined, independent of how it is parameterized. The electromagnetic field, on the other hand, are defined on the space-time. The geometric Lagrangian density of the system will be a function of the 4-potential of electromagnetic fields and also a functional of particles' world-lines. This is qualitatively different from the standard field theory, where the Lagrangian density is a local function of field [8]. The other difficulty associated with the geometric setting is due to the mass-shell condition, which exist even for the geometric variational principle for a single particle [9]. The standard EL equation will be generalized into the geometric EL equation when the mass-shell condition is imposed. For the particle-field system, the geometric EL equation is further generalized into a weak geometric EL equation.

This paper is organized as follows. In Sec. II, we will discuss how to expand Qin's work into the relativistic situation in a nongeometric way. The geometric Lagrangian for a single particle and the corresponding geometric EL equation are discussed in Sec. III. In Sec. IV, the geometric Lagranian density for particle-field system and its properties are discussed. The geometric weak EL equation and the link it provides between conservation laws and space-time symmetries are given in Sec. V.

II. NONGEOMETRIC FIELD THEORY AND WEAK EULER-LAGRANGE EQUATION FOR RELATIVISTIC PARTICLE-FIELD SYSTEMS

The classical relativistic particle-field syetems in flat space is governed by the following Newton-Maxwell equations,

$$\frac{d}{dt}(\gamma_{sp} m_s \dot{\mathbf{X}}_{sp}) = q_s(\mathbf{E}+\frac{1}{c}\dot{\mathbf{X}}_{sp}\times\mathbf{B}), \tag{12}$$

$$\nabla\cdot\mathbf{E} = 4\pi\sum_{s,p} q_s\delta(\mathbf{X}_{sp}-\boldsymbol{x}), \tag{13}$$

$$\nabla\times\mathbf{B} = \frac{4\pi}{c}\sum_{s,p} q_s\dot{\mathbf{X}}_{sp}\delta(\mathbf{X}_{sp}-\boldsymbol{x}) + \frac{1}{c}\frac{\partial\mathbf{E}}{\partial t}, \tag{14}$$

$$\nabla\times\mathbf{E} = -\frac{1}{c}\frac{\partial\mathbf{B}}{\partial t}, \tag{15}$$

$$\nabla \cdot \mathbf{B} = 0, \tag{16}$$

where $\gamma_{sp} \equiv 1/\sqrt{1 - \dot{\boldsymbol{X}}_{sp}^2/c^2}$ is the relativistic factor for p-th particle of s-species, m_s is the mass for any particle of s-species, $\mathbf{X}_{sp}$ is the trajectory, $\boldsymbol{x}$ is a point in the configuration space, and $\delta(\mathbf{X}_{sp} - \boldsymbol{x})$ is the Dirac delta function on 3D configuration space [10]. The electric field $\mathbf{E}(t, \boldsymbol{x})$ and magnetic field $\mathbf{B}(t, \boldsymbol{x})$ are functions of space and time. For this system, the action $\mathcal{A}$ is

$$\mathcal{A}[\varphi, \mathbf{A}, \mathbf{X}_{sp}] = \int \mathcal{L}_{PF} dt d^3x, \tag{17}$$

where

$$\begin{aligned}\mathcal{L}_{PF} = &- \sum_{s,p} \gamma_{sp}^{-1} m_s c^2 \delta(\mathbf{X}_{sp} - \boldsymbol{x}) + \frac{q_s}{c} \mathbf{A} \cdot \dot{\mathbf{X}}_{sp} \delta(\mathbf{X}_{sp} - \boldsymbol{x}) - q_s \varphi \delta(\mathbf{X}_{sp} - \boldsymbol{x}) \\ &+ \frac{1}{8\pi} \left[\left(-\frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} - \nabla \varphi \right)^2 - \nabla \times \boldsymbol{A} \right]\end{aligned} \tag{18}$$

is the Lagrangian density of this system [6]. The variation of $\mathcal{A}$ induced by $\delta \mathbf{X}_{sp}$, $\delta \mathbf{A}$ and $\delta \varphi$ is

$$\delta \mathcal{A} = \sum_{s,p} \int dt \delta \mathbf{X}_{sp} \cdot \int E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF}) d^3x + \int E_{\varphi}(\mathcal{L}_{PF}) \delta \varphi dt d^3x + \int E_{\mathbf{A}}(\mathcal{L}_{PF}) \cdot \delta \mathbf{A} dt d^3x, \tag{19}$$

where

$$E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF}) \equiv \frac{\partial \mathcal{L}_{PF}}{\partial \mathbf{X}_{sp}} - \frac{D}{Dt} \cdot \frac{\partial \mathcal{L}_{PF}}{\partial \dot{\mathbf{X}}_{sp}}, \tag{20}$$

$$E_{\varphi}(\mathcal{L}_{PF}) \equiv \frac{\partial \mathcal{L}_{PF}}{\partial \varphi} - \frac{D}{D\boldsymbol{x}} \cdot \frac{\partial \mathcal{L}_{PF}}{\partial \dot{\varphi}}, \tag{21}$$

$$E_{\mathbf{A}}(\mathcal{L}_{PF}) \equiv \frac{\partial \mathcal{L}_{PF}}{\partial \mathbf{A}} - \frac{D}{D\boldsymbol{x}} \cdot \frac{\partial \mathcal{L}_{PF}}{\partial \nabla \mathbf{A}} - \frac{D}{Dt} \frac{\partial \mathcal{L}_{PF}}{\partial \mathbf{A}_{,t}}. \tag{22}$$

For $\delta \mathcal{A} = 0$, we have

$$\int E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF}) d^3\boldsymbol{x} = 0, \tag{23}$$

$$E_{\varphi}(\mathcal{L}_{PF}) = 0, \tag{24}$$

$$E_{\mathbf{A}}(\mathcal{L}_{PF}) = 0 \tag{25}$$

due to the arbitrarinesses of $\delta \mathbf{X_{sp}}$, $\delta \varphi$ and $\delta \mathbf{A}$ in Eq. (19). Equation (23) will be called submanifold EL equation because it is defined only on the time-axis after the integrating

over spatial variable. Moreover, substituting Eq. (18) into Eqs. (20)-(25), we will recover Eqs. (12)-(16) by defining $\mathbf{E} = -\partial\mathbf{A}/\partial t - \nabla\varphi$ and $\mathbf{B} = \nabla \times \mathbf{A}$.

For Eq. (23), in general, we expect that $E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF}) \neq 0$ although the integral of this term vanishes. We can derive an explicit expression for $E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF})$ as

$$E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF}) = \frac{\partial \mathcal{L}_{PF}}{\partial \mathbf{X}_{sp}} - \frac{D}{Dt} \cdot \frac{\partial \mathcal{L}_{PF}}{\partial \dot{\mathbf{X}}_{sp}} = \frac{\partial}{\partial x}\left(H_{sp} - \mathbf{P}_{sp} \cdot \dot{\mathbf{X}}_{sp}\right) + \frac{\partial}{\partial \boldsymbol{x}} \cdot \left(\dot{\mathbf{X}}_{sp}\mathbf{P}_{sp}\right), \tag{26}$$

where

$$H_{sp} = \left(\gamma_{sp} m_s c^2 + q_s \varphi\right) \delta\left(\mathbf{X}_{sp} - \boldsymbol{x}\right), \tag{27}$$

$$\mathbf{P}_{sp} = \left(\gamma_{sp} m_s \dot{\mathbf{X}}_{sp} + q_s \mathbf{A}\right) \delta\left(\mathbf{X}_{sp} - \boldsymbol{x}\right). \tag{28}$$

Equation (26) is called weak EL equation [7], and the qualifer "weak" is used to indicate the fact that only the spatial integral of $E_{\mathbf{X_{sp}}}(\mathcal{L}_{PF})$ is zero (see Eq. (23)).

Next, we define the symmetry of the action $\mathcal{A}[\varphi, \mathbf{A}, \mathbf{X}_{sp}]$ to be a group of transformation

$$(t, \boldsymbol{x}, \varphi, \mathbf{A}, \mathbf{X}_{sp}) \longmapsto \left(\tilde{t}, \tilde{\boldsymbol{x}}, \tilde{\varphi}, \tilde{\mathbf{A}}, \tilde{\mathbf{X}}_{sp}\right), \tag{29}$$

such that

$$\int \mathcal{L}_{PF}\left(t, \boldsymbol{x}, \varphi, \mathbf{A}, \mathbf{X}_{sp}\right) dt d^3x = \int \tilde{\mathcal{L}}_{PF}\left(\tilde{t}, \tilde{\boldsymbol{x}}, \tilde{\varphi}, \tilde{\mathbf{A}}, \tilde{\mathbf{X}}_{sp}\right) dt d^3\boldsymbol{x}. \tag{30}$$

For our Lagrangian density (see Eq. (18)), if the group transformation is the time-translation,

$$\left(\tilde{t}, \tilde{\boldsymbol{x}}, \tilde{\varphi}, \tilde{\mathbf{A}}, \tilde{\mathbf{X}}_{sp}\right) = \left(t + \epsilon, \boldsymbol{x}, \varphi, \mathbf{A}, \mathbf{X}_{sp}\right), \quad \epsilon \in \mathbb{R} \tag{31}$$

the condition (30) will be satisfied because

$$\frac{\partial \mathcal{L}_{PF}}{\partial t} = 0. \tag{32}$$

Using weak EL equations (26) for particles, the EL equations for fields (see Eqs. (21), (22), (24), (25)), we obtain the energy conservation law

$$\frac{\partial}{\partial t}\left[\sum_{s,p} \gamma_{sp} m_s c^2 \delta\left(\mathbf{X}_{sp} - \boldsymbol{x}\right) + \frac{\mathbf{E^2} + \mathbf{B^2}}{8\pi}\right] + \nabla \cdot \left[\sum_{s,p} \gamma_{sp} m_s c^2 \dot{\mathbf{X}}_{sp} \delta\left(\mathbf{X}_{sp} - \boldsymbol{x}\right) + \frac{c}{4\pi}\mathbf{E} \times \mathbf{B}\right] = 0. \tag{33}$$

Equation (30) holds for spatial-translation as well

$$\left(\tilde{t}, \tilde{\boldsymbol{x}}, \tilde{\varphi}, \tilde{\mathbf{A}}, \tilde{\mathbf{X}}_{sp}\right) = \left(t, \boldsymbol{x} + \epsilon\mathbf{X}, \varphi, \mathbf{A}, \mathbf{X}_{sp}\epsilon\mathbf{X}\right), \quad \epsilon \in \mathbb{R} \tag{34}$$

because $\mathcal{L}_{PF}$ satisfies

$$\frac{\partial \mathcal{L}_{PF}}{\partial \boldsymbol{x}} + \sum_{s,p} \frac{\partial \mathcal{L}_{PF}}{\partial \mathbf{X}_{sp}} = 0. \tag{35}$$

As a consequence, the momentum conservation law due to this symmetry can be written as

$$\frac{\partial}{\partial t}\left[\sum_{s,p} \gamma_{sp} m_S \dot{\mathbf{X}}_{sp} + \frac{1}{4\pi c}\mathbf{E}\times\mathbf{B}\right] + \nabla\cdot\left[\sum_{s,p} \gamma_{sp} m_S \dot{\mathbf{X}}_{sp}\dot{\mathbf{X}}_{sp} + \frac{\mathbf{E^2}+\mathbf{B^2}}{8\pi}\mathbf{I} - \frac{\mathbf{EE}+\mathbf{BB}}{4\pi}\right] = 0. \tag{36}$$

Details of the derivation can be found in Ref. [7]. However, in relativistic situation, this split form of space and time is not elegant. The space and time should be treated as one identity in the most fundamental and geometric approach. In the following sections, we will explore the geometric ways to establish the relations between symmetries and conservation laws.

III. GEOMETRIC LAGRANGIAN AND GEOMETRIC EULER-LAGRANGIAN EQUATION FOR A SINGLE PARTICLE

For a classical particle, the action can be expressed as

$$\mathcal{A} = \int_{t_1}^{t_2} L(\boldsymbol{x}, \dot{\boldsymbol{x}}, t) dt, \tag{37}$$

where $L(\boldsymbol{x}, \dot{\boldsymbol{x}}, t)$ is the Lagrangian[8, 11]. Apply the principle of least action, $\delta\mathcal{A} = 0$, we have

$$0 = \delta\mathcal{A} = \int_{t_1}^{t_2} \delta L(\boldsymbol{x}, \dot{\boldsymbol{x}}, t) dt = \int_{t_1}^{t_2} \left[\frac{\partial L}{\partial \boldsymbol{x}} - \frac{d}{dt}\left(\frac{\partial L}{\partial \dot{\boldsymbol{x}}}\right)\right]\cdot \delta\boldsymbol{x} dt + \left[\frac{\partial L}{\partial \dot{\boldsymbol{x}}}\cdot\delta\boldsymbol{x}\right]_{t_1}^{t_2}. \tag{38}$$

Because of the fact that $\delta\boldsymbol{x}(t_1) = \delta\boldsymbol{x}(t_2) = 0$ and the arbitrariness of $\delta\boldsymbol{x}$, we obtain the EL equation for the dynamics of the particle

$$\frac{\partial L}{\partial \boldsymbol{x}} - \frac{d}{dt}\left(\frac{\partial L}{\partial \dot{\boldsymbol{x}}}\right) = 0. \tag{39}$$

For relativistic particles, the variational principle is more involved. In special relativity, the Lagrangian should be Lorentz invariant [12]. For this reason, the integral variable of the action should be the proper time τ and the action is

$$\mathcal{A} = \int_{p_1}^{p_2} \widetilde{L}(\chi, \dot{\chi}, \tau) d\tau, \tag{40}$$

where χ and $\dot{\chi} \equiv d\chi/d\tau$ are the world-line and 4-velocity of the particle, respectively. The Lagrangian $\widetilde{L}$ is manifestly covariant [12] and is called geometric Lagrangian in the present study. The integral in Eq. (40) is along any possible world-lines passing p_1 and p_2 in space-time. If we parameterize the world-line by the proper time τ, then the Eq. (40) can be written as

$$\mathcal{A} = \int_{\tau_1}^{\tau_2} \widetilde{L}(\chi, \dot{\chi}, \tau) d\tau, \tag{41}$$

where τ_1 is the proper time at the beginning and τ_2 is the proper time at the end. We can choose the parameter τ_1 to be same for any possible world-line. But, the parameter τ_2 are generally different because of the different length for different world-lines passing the given space-time point p_2, which means $\delta(d\tau) \neq 0$. This is different from the non-relativistic case, which implicitly assumes $\delta(dt) = 0$. The Hamiltonian principle is

$$0 = \delta\mathcal{A} = \int_{\tau_1}^{\tau_2} \delta\widetilde{L}(\chi, \dot{\chi}, \tau) d\tau + \int_{\tau_1}^{\tau_2} \widetilde{L}(\chi, \dot{\chi}, \tau) \delta(d\tau), \tag{42}$$

where

$$\begin{aligned} \delta(d\tau) &= \frac{1}{c^2}\delta\left(\frac{d\chi^\mu d\chi_\mu}{d\tau}\right) = \frac{1}{c^2}\delta\left[d\chi^\mu \frac{d\chi_\mu}{d\tau}\right] = \frac{1}{c^2}\delta\left[d\chi^\mu \dot{\chi}_\mu\right] \\ &= \frac{1}{c^2}\delta\left(d\chi^\mu\right)\dot{\chi}_\mu + \frac{1}{c^2} d\chi^\mu \delta\left(\dot{\chi}_\mu\right) = \frac{1}{c^2}\delta\left(d\chi^\mu\right)\dot{\chi}_\mu + \frac{1}{c^2}\dot{\chi}^\mu \delta\left(\dot{\chi}_\mu\right) d\tau. \end{aligned} \tag{43}$$

Here, χ^μ and $\dot{\chi}^\mu$ ($\mu = 0, 1, 2, 3$) are the components of the world-line and 4-veolocity of the particle. From the mass-shell condition

$$P^\mu P_\mu = m_0^2 c^2 \text{ or } \dot{\chi}^\mu \dot{\chi}_\mu = c^2, \tag{44}$$

where $P^\mu = m_0 \dot{\chi}^\mu$ are the components of 4-momentum and m_0 is the rest mass,

$$0 = \delta\left(c^2\right) = \delta\left(\dot{\chi}^\mu \dot{\chi}_\mu\right) = \delta\left(\dot{\chi}^\mu\right)\dot{\chi}_\mu + \dot{\chi}^\mu \delta\left(\dot{\chi}_\mu\right) = 2\dot{\chi}^\mu \delta\left(\dot{\chi}_\mu\right), \tag{45}$$

which means that Eq. (43) can be reduced to

$$\delta(d\tau) = \frac{1}{c^2}\dot{\chi}_\mu \delta\left(d\chi^\mu\right). \tag{46}$$

The second term on the right-hand side of Eq. (42) is

$$\begin{aligned}\frac{1}{c^2}\int_{\tau_1}^{\tau_2}\widetilde{L}\dot{\chi}_\mu\delta\left(d\chi^\mu\right) &= \frac{1}{c^2}\int_{\tau_1}^{\tau_2}\widetilde{L}\dot{\chi}_\mu\frac{d\left(\delta\chi^\mu\right)}{d\tau}d\tau \\ &= -\frac{1}{c^2}\int_{\tau_1}^{\tau_2}\frac{d}{d\tau}\left[\widetilde{L}\dot{\chi}_\mu\right]\left(\delta\chi^\mu\right)d\tau + \left[\widetilde{L}\dot{\chi}_\mu\left(\delta\chi^\mu\right)\right]_{\tau_1}^{\tau_2} \\ &= -\frac{1}{c^2}\int_{\tau_1}^{\tau_2}\frac{d}{d\tau}\left[\widetilde{L}\dot{\chi}_\mu\right]\delta\chi^\mu d\tau. \end{aligned} \tag{47}$$

The first term on the right-hand side of Eq. (42) is

$$\begin{aligned}&\int_{\tau_1}^{\tau_2}\left[\frac{\partial\widetilde{L}}{\partial\chi^\mu}\delta\chi^\mu + \frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu}\delta\dot{\chi}^\mu\right]d\tau \\ &= \int_{\tau_1}^{\tau_2}\left\{\frac{\partial\widetilde{L}}{\partial\chi^\mu}\delta\chi^\mu + \frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu}\left[\frac{d\left(\delta\chi^\mu\right)}{d\tau} - \frac{1}{c^2}\dot{\chi}^\mu\dot{\chi}_\nu\frac{d\left(\delta\dot{\chi}^\nu\right)}{d\tau}\right]\right\}d\tau \qquad (48) \\ &= \int_{\tau_1}^{\tau_2}\frac{\partial\widetilde{L}}{\partial\chi^\mu}\delta\chi^\mu d\tau + \int_{\tau_1}^{\tau_2}\left[\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu}\dot{\chi}^\nu\dot{\chi}_\mu\right]\frac{d\left(\delta\chi^\mu\right)}{d\tau}d\tau \\ &= \int_{\tau_1}^{\tau_2}\frac{\partial\widetilde{L}}{\partial\chi^\mu}\delta\chi^\mu d\tau - \int_{\tau_1}^{\tau_2}\frac{d}{d\tau}\left[\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\dot{\chi}^\nu\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu}\dot{\chi}_\mu\right]\left(\delta\chi^\mu\right)d\tau \\ &+ \left\{\left[\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\dot{\chi}^\nu\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu}\dot{\chi}_\mu\right]\delta\chi^\mu\right\}_{\tau_1}^{\tau_2} \\ &= \int_{\tau_1}^{\tau_2}\left[\frac{\partial\widetilde{L}}{\partial\chi^\mu} - \frac{d}{d\tau}\left(\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\dot{\chi}^\nu\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu}\dot{\chi}_\mu\right)\right]\delta\chi^\mu d\tau, \qquad (49)\end{aligned}$$

where used if made of the following identity,

$$\delta\dot{\chi}^\mu = \frac{d\left(\delta\chi^\mu\right)d\tau - \frac{1}{c^2}\chi_\nu d\chi^\mu d\left(\delta\chi^\nu\right)}{\left(d\tau\right)^2} = \frac{d\left(\delta\chi^\mu\right)}{d\tau} - \frac{1}{c^2}\dot{\chi}^\mu\dot{\chi}_\nu\frac{d\left(\delta\chi^\nu\right)}{d\tau}. \tag{50}$$

Substituting Eqs. (47) and (49) into Eq. (42), we obtain

$$0 = \delta\mathcal{A} = \int_{\tau_1}^{\tau_2}\left\{\frac{\partial\widetilde{L}}{\partial\chi^\mu} - \frac{d}{d\tau}\left[\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\left(\dot{\chi}^\nu\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu} - \widetilde{L}\right)\dot{\chi}_\mu\right]\right\}\delta\chi^\mu d\tau,$$

which implies

$$\frac{\partial\widetilde{L}}{\partial\chi^\mu} - \frac{d}{d\tau}\left[\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\mu} - \frac{1}{c^2}\left(\dot{\chi}^\nu\frac{\partial\widetilde{L}}{\partial\dot{\chi}^\nu} - \widetilde{L}\right)\dot{\chi}_\mu\right] = 0. \tag{51}$$

due to the arbitrariness of $\delta\chi^\mu$. Equation (51) will be called geometric EL equation. We note that it has been derived by Barut using the method of Lagrange multiplier [9].

IV. GEOMETRIC LAGRANGIAN DENSITY

For particle-field systems, we need to find the density of the geometric Lagrangian. We start from Eq. (17), the nongeometric form of the action of our system. It can be written as [6]

$$\mathcal{A} = -\sum_{s,p} \int m_s c^2 d\tau - \sum_{s,p} \int \frac{q_s}{c} A_\mu d\chi^\mu_{sp} - \frac{1}{16\pi c} \int F_{\mu\nu} F^{\mu\nu} d\Omega, \tag{52}$$

where χ^μ_{sp} ($\mu = 0, 1, 2, 3$) are components of the world-line of pth particle of s-species, τ is the proper time, $d\Omega$ is the 4 dimensional volume element in Minkowshi space-time, and the 4-potential A_μ and the field-strength tensor $F^{\mu\nu}$ are the functions on the Minkowskian space. The boundary of the domain is taken to be at the infinity.

Equation (52) can be easily translated into another form,

$$\mathcal{A} = \sum_{s,p} \int \left(-m_s c^2 - \frac{q_s}{c} A_\mu \dot{\chi}^\mu(\tau) \right) d\tau - \frac{1}{4\pi c} \int \partial^{[\mu} A^{\nu]} \partial_\mu A_\nu d\Omega, \tag{53}$$

by using the relations $\dot{\chi}^\mu = d\chi^\mu/d\tau$ and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, where the symbol $[\mu\ \nu]$ in Eq. (53) means the antisymmetrization of the indexes of μ and ν [5]. The action is manifestly covariant or geometric. However, to obtain the Lagrangian density of the particle-field systems, we should change it into another form by multiplying the first term in Eq. (53) with the equation

$$\int \delta\left(\chi_{sp} - \chi\right) d\Omega = 1, \tag{54}$$

where χ is an arbitrary world point in Minkowskian space, and $\delta\left(\chi_{sp} - \chi\right)$ is the Dirac delta function [10]. Then equation (53) becomes

$$\mathcal{A} = \int \left[\sum_{s,p} \int \left(-m_s c^2 - \frac{q_s}{c} A_\mu \dot{\chi}^\mu(\tau) \right) \delta\left(\chi_{sp} - \chi\right) d\tau \right] d\Omega - \frac{1}{4\pi c} \int \partial^{[\mu} A^{\nu]} \partial_\mu A_\nu d\Omega. \tag{55}$$

The geometric Lagrangian density is easy to read off from Eq. (55) as

$$\mathcal{L} = \sum_{s,p} \int \left(-m_s c^2 - \frac{q_s}{c} A_\mu \dot{\chi}^\mu_{sp}(\tau) \right) \delta\left(\chi_{sp} - \chi\right) d\tau - \frac{1}{4\pi c} \partial^{[\mu} A^{\nu]} \partial_\mu A_\nu. \tag{56}$$

The geometric Lagrangian above can be simplified as

$$\mathcal{L} = \mathcal{L}_P + \mathcal{L}_F = \int \hat{\mathcal{L}}_P d\tau + \mathcal{L}_F \tag{57}$$

by defining

$$\hat{\mathcal{L}}_P = \sum_{s,p} \left(-m_s c^2 - \frac{q_s}{c} A_\mu \dot{\chi}^\mu_{sp}(\tau) \right) \delta \left(\chi_{sp} - \chi \right), \tag{58}$$

$$\mathcal{L}_P = \int \hat{\mathcal{L}}_P \left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A \right) d\tau \tag{59}$$

and

$$\mathcal{L}_F = -\frac{1}{4\pi c} \partial^{[\mu} A^{\nu]} \partial_\mu A_\nu. \tag{60}$$

Here, $\hat{\mathcal{L}}_P$ is a function of χ , χ_{sp} , $\dot{\chi}_{sp}$ and A. On the other hand, $\mathcal{L}_P$ can be regarded as a functional of the world-lines of particles and a function of the space-time and the field A.

V. GEOMETRIC WEAK EULER-LAGRANGIAN EQUATION AND ENERGY-MOMENTUM CONSERVATION

Now we determine how the action given by Eq. (55) and geometric Lagrangian density vary in response to the variation of $\delta\chi_{sp}$ and δA. From Eqs. (55)-(60), the variation of action of the particle-field system can be written as

$$\begin{aligned} \delta\mathcal{A} = & \int \delta \left[\int \hat{\mathcal{L}}_P \left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A \right) d\tau \right]_A d\Omega \\ & + \int \delta \left[\int \hat{\mathcal{L}}_P \left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A \right) d\tau \right]_{\chi_{sp}, \dot{\chi}_{sp}} d\Omega + \int \delta \mathcal{L}_F d\Omega, \end{aligned} \tag{61}$$

where the notation $\delta \left[\int \hat{\mathcal{L}}_P \left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A \right) d\tau \right]_\alpha$ $(\alpha = A, \chi_{sp}, \dot{\chi}_{sp})$ means keeping α fixed.

For the first term on the right-hand side of Eq. (61), it can be treated by the same procedure for the derivation of Eq. (51),

$$\begin{aligned} & \int \delta \left[\int \hat{\mathcal{L}}_P \left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A \right) d\tau \right]_A d\Omega \\ & = \int \int \left\{ \frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^\mu_{sp}} - \frac{D}{D\tau} \left[\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\mu_{sp}} - \frac{1}{c^2} \left(\chi^\nu_{sp} \frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\nu_{sp}} - \hat{\mathcal{L}}_P \right) \dot{\chi}_{sp\mu} \right] \right\} \delta\chi^\mu_{sp} d\tau d\Omega \\ & = \int \delta\chi^\mu_{sp} d\tau \int \left\{ \frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^\mu_{sp}} - \frac{D}{D\tau} \left[\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\mu_{sp}} - \frac{1}{c^2} \left(\chi^\nu_{sp} \frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\nu_{sp}} - \hat{\mathcal{L}}_P \right) \dot{\chi}_{sp\mu} \right] \right\} d\Omega. \end{aligned} \tag{62}$$

The second and third terms on the right-hand side of Eq. (61) are actually

$$\int \left\{ \frac{\partial \mathcal{L}}{\partial A_\mu} - \frac{D}{D\chi^\nu} \left[\frac{\partial \mathcal{L}}{\partial \left(\partial_\nu A_\mu \right)} \right] \right\} \delta A_\mu d\Omega. \tag{63}$$

If we define

$$E_A = \frac{\partial \mathcal{L}}{\partial A_\mu} - \frac{D}{D\chi^\nu}\left[\frac{\partial \mathcal{L}}{\partial\left(\partial_\nu A_\mu\right)}\right], \tag{64}$$

$$E_{\chi_{sp}} = \frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^\mu_{sp}} - \frac{D}{D\tau}\left[\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\mu_{sp}} - \frac{1}{c^2}\left(\chi^\nu_{sp}\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\nu_{sp}} - \hat{\mathcal{L}}_P\right)\dot{\chi}_{sp\mu}\right], \tag{65}$$

and substitute Eqs. (62)-(65) into Eq. (61), then

$$\delta\mathcal{A} = \int\left[\int E_{\chi_{sp}} d\Omega\right]\delta\chi^\mu_{sp} d\tau + \int E_A \delta A_\mu d\Omega. \tag{66}$$

Thus,

$$E_A = \frac{\partial \mathcal{L}}{\partial A_\mu} - \frac{D}{D\chi^\nu}\left[\frac{\partial \mathcal{L}}{\partial\left(\partial_\nu A_\mu\right)}\right] = 0, \tag{67}$$

$$\int E_{\chi_{sp}} d\Omega \equiv \int\left\{\frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^\mu_{sp}} - \frac{D}{D\tau}\left[\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\mu_{sp}} - \frac{1}{c^2}\left(\chi^\nu_{sp}\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^\nu_{sp}} - \hat{\mathcal{L}}_P\right)\dot{\chi}_{sp\mu}\right]\right\} d\Omega = 0, \tag{68}$$

by the arbitrariness of $\delta\chi^\mu_{sp}$ and δA_μ.

Equation (67) is the EL equation of the 4-potential of the electromagnetic field. Substituting Eq. (56) into Eq. (67) gives the Maxwell equation,

$$\partial^\nu\left(\partial_\nu A_\mu - \partial_\mu A_\nu\right) = -\frac{4\pi}{c}\sum_{s,p} q_s \int \dot{\chi}_{sp\mu}\delta\left(\chi_{sp} - \chi\right) ds, \tag{69}$$

where s is the length of the world-line, i.e.,

$$ds = cd\tau. \tag{70}$$

The 4-current [13] is

$$J_\mu = \sum_{s,p} q_s \int \dot{\chi}_{sp\mu}\delta\left(\chi_{sp} - \chi\right) ds. \tag{71}$$

Equation (68), will be called geometric submanifold EL equation because it is defined only on the world-line after the integrating over the space-time variable [7]. If χ_{sp} were a function of the entire space-time domain, then $E_{\chi_{sp}}$ would vanish everywhere, as in the case for 4-potential A. Generally, we expect that $E_{\chi_{sp}} \neq 0$.

We now derive an expression for $E_{\chi_{sp}}$ by substituting Eq. (58) into Eq. (65). For the first term in $E_{\chi_{sp}}$,

$$\begin{aligned}\frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^\mu_{sp}} &= (-m_s c^2 - \frac{q_s}{c}A_\nu \dot{\chi}^\nu_{sp})\frac{\partial \delta_2}{\partial \chi^\mu_{sp}} \\ &= \frac{D}{D\chi^\mu}\left[\left(m_s c^2 + \frac{q_s}{c}A_\nu\dot{\chi}^\nu_{sp}\right)\delta_2\right] - \frac{q_s}{c}\frac{\partial A_\nu}{\partial \chi^\mu}\dot{\chi}^\nu_{sp}\delta_2,\end{aligned} \tag{72}$$

where $\delta_2 \equiv \delta\left(\chi_{sp} - \chi\right)$. The second term of $E_{\chi_{sp}}$ is given by

$$\begin{aligned} -\frac{D}{D\tau}\left(\frac{\partial\hat{\mathcal{L}}_P}{\partial\dot{\chi}^{\mu}_{sp}}\right) &= \frac{q_s}{c}A_\mu\frac{D\delta_2}{D\tau} = -\frac{q_s}{c}A_\mu\dot{\chi}^{\nu}_{sp}\frac{\partial\delta_2}{\partial\chi^{\nu}} \\ &= -\frac{D}{D\chi^{\nu}}\left[\frac{q_s}{c}A_\mu\dot{\chi}^{\nu}_{sp}\delta_2\right] + \frac{q_s}{c}\dot{\chi}^{\nu}_{sp}\frac{\partial A_\mu}{\partial\chi^{\nu}}\delta_2 \end{aligned} \tag{73}$$

The third and fourth terms came from the mass-shell condition, and can be rewritten as

$$\begin{aligned} &-\frac{D}{D\tau}\left(-\frac{1}{c^2}\dot{\chi}^{\nu}_{sp}\frac{\partial\hat{\mathcal{L}}_P}{\partial\dot{\chi}^{\nu}_{sp}}\dot{\chi}_{sp\mu}\right) \\ &= -\frac{q_s}{c^3}A_\nu\ddot{\chi}^{\nu}_{sp}\dot{\chi}_{sp\mu}\delta_2 - \frac{q_s}{c^3}A_\nu\dot{\chi}^{\nu}_{sp}\ddot{\chi}_{sp\mu}\delta_2 - \frac{q_s}{c^3}A_\nu\dot{\chi}^{\nu}_{sp}\dot{\chi}_{sp\mu}\frac{D\delta_2}{D\tau} \\ &= -\frac{q_s}{c^3}A_\nu\ddot{\chi}^{\nu}_{sp}\dot{\chi}_{sp\mu}\delta_2 - \frac{q_s}{c^3}A_\nu\dot{\chi}^{\nu}_{sp}\ddot{\chi}_{sp\mu}\delta_2 + \frac{D}{D\chi^{\sigma}}\left(\frac{q_s}{c^3}A_\nu\dot{\chi}^{\nu}_{sp}\dot{\chi}_{sp\mu}\dot{\chi}^{\sigma}_{sp}\delta_2\right) \\ &-\frac{q_s}{c^3}\dot{\chi}^{\nu}_{sp}\frac{\partial A_\nu}{\partial\chi^{\sigma}}\dot{\chi}^{\sigma}_{sp}\dot{\chi}_{sp\mu}\delta_2, \end{aligned} \tag{74}$$

and

$$\begin{aligned} &-\frac{D}{D\tau}\left(\frac{1}{c^2}\hat{\mathcal{L}}_P\dot{\chi}_{sp\mu}\right) \\ &= -\frac{D}{D\chi^{\nu}}\left[\left(m_s + \frac{q_s}{c^3}A_\sigma\dot{\chi}^{\sigma}_{sp}\right)\dot{\chi}_{sp\mu}\dot{\chi}^{\nu}_{sp}\delta_2\right] + \frac{q_s}{c^3}\dot{\chi}^{\nu}_{sp}\frac{\partial A_\sigma}{\partial\chi^{\nu}}\dot{\chi}^{\sigma}_{sp}\dot{\chi}_{sp\mu}\delta_2 \\ &+m_s\ddot{\chi}_{sp\mu}\delta_2 + \frac{q_s}{c^3}A_\sigma\dot{\chi}^{\sigma}_{sp}\ddot{\chi}_{sp\mu}\delta_2 + \frac{q_s}{c^3}A_\sigma\ddot{\chi}^{\sigma}_{sp}\dot{\chi}_{sp\mu}\delta_2. \end{aligned} \tag{75}$$

Therefore,

$$\begin{aligned} E_{\chi_{sp}} = \frac{D}{D\chi^{\nu}}&\left\{\left[\left(m_sc^2 + \frac{q_s}{c}A_\sigma\dot{\chi}^{\sigma}_{sp}\right)\eta_{\mu}^{\ \nu} - \left(\frac{q_s}{c}A_\mu + m_s\dot{\chi}_{sp\mu}\right)\dot{\chi}^{\nu}_{sp}\right]\delta_2\right\} \\ &+\left[m_s\ddot{\chi}_{sp\mu} - \frac{q_s}{c}\dot{\chi}^{\nu}_{sp}\left(\frac{\partial A_\nu}{\partial\chi^{\mu}} - \frac{\partial A_\mu}{\partial\chi^{\nu}}\right)\right]\delta_2. \end{aligned} \tag{76}$$

Substituting Eq. (76) into the geometric submanifold EL equation (68), we immediately obtain the equation of motion for particles,

$$m_s\ddot{\chi}_{sp\mu} = \frac{q_s}{c}\left(\frac{\partial A_\nu}{\partial\chi^{\mu}} - \frac{\partial A_\mu}{\partial\chi^{\nu}}\right)\dot{\chi}^{\nu}_{sp}. \tag{77}$$

The right term in Eq. (77) is the 4-Lorenzen force on particles. Then, $E_{\chi_{sp}}$ reduces to

$$\begin{aligned} E_{\chi_{sp}} &= \frac{\partial\hat{\mathcal{L}}_P}{\partial\chi^{\mu}_{sp}} - \frac{D}{D\tau}\left[\frac{\partial\hat{\mathcal{L}}_P}{\partial\dot{\chi}^{\mu}_{sp}} - \frac{1}{c^2}\left(\dot{\chi}^{\nu}_{sp}\frac{\partial\hat{\mathcal{L}}_P}{\partial\dot{\chi}^{\nu}_{sp}} - \hat{\mathcal{L}}_P\right)\dot{\chi}_{sp\mu}\right] \\ &= \frac{D}{D\chi^{\nu}}\left\{\left[\left(m_sc^2 + \frac{q_s}{c}A_\sigma\dot{\chi}^{\sigma}_{sp}\right)\eta_{\mu}^{\ \nu} - \left(\frac{q_s}{c}A_\mu + m_s\dot{\chi}_{sp\mu}\right)\dot{\chi}^{\nu}_{sp}\right]\delta_2\right\}. \end{aligned} \tag{78}$$

As expected, $E_{\chi_{sp}} \neq 0$. We will refer to Eq. (78) as the geometric weak EL equation [7], which play a significant role in subsequent analysis of the local conservation laws. The

qualifier "weak" is used to indicate the fact that only the space-time integral of $E_{\chi_{sp}}$ is zero. Unlike the non-relativistic case (see Eq. (26)), the geometric weak EL equation here is written as a differential equation about $\hat{\mathcal{L}_P}$, instead of the Lagrangian density.

Another crucial step to systematically obtain conservation laws is finding the symmetries. A symmetry group of action $\mathcal{A}$ is defined by a continuous transformation

$$(\chi, [\chi_{sp}], [\dot{\chi}_{sp}], A, \partial A) \longmapsto \left(\widetilde{\chi}, [\widetilde{\chi}_{sp}], [\widetilde{\dot{\chi}}_{sp}], \widetilde{A}, \widetilde{\partial A}\right) \tag{79}$$

such that

$$\int \mathcal{L}\left(\chi, [\chi_{sp}], [\dot{\chi}_{sp}], A, \partial A\right) d\Omega = \int \tilde{\mathcal{L}}\left(\widetilde{\chi}, [\widetilde{\chi}_{sp}], [\widetilde{\dot{\chi}}_{sp}], \widetilde{A}, \widetilde{\partial A}\right) d\Omega, \tag{80}$$

where the indexes for physical quantities has been omitted to simplify the notations. The symbol $[\beta]\left(\beta = \chi_{sp}, \dot{\chi}_{sp}, \widetilde{\chi}_{sp}, \widetilde{\dot{\chi}}_{sp}\right)$ indicates that the geometric Lagrangian density $\mathcal{L}$ is a functional of β. For the following transformation defined by

$$\left(\widetilde{\chi}, \widetilde{\chi}_{sp}, \widetilde{\dot{\chi}}_{sp}, \widetilde{A}, \widetilde{\partial A}\right) = \left(\chi + \epsilon X, \chi_{sp} + \epsilon X, \dot{\chi}_{sp}, A, \partial A\right), \quad \epsilon \in R \tag{81}$$

where X is a given constant 4-vector field in Minkowsikian space, the condition (80) is satisfied because

$$\mathcal{L} = \int \hat{\mathcal{L}_P}\left(\chi_{sp}, \dot{\chi}_{sp}, \chi, A\right) d\tau + \mathcal{L}_F = \int \hat{\mathcal{L}_P}\left(\chi + \epsilon X, \chi_{sp} + \epsilon X, \dot{\chi}_{sp}, A\right) d\tau + \mathcal{L}_F, \tag{82}$$

with $\widetilde{A}(\widetilde{\chi}) = A(\chi) = A(\widetilde{\chi} - \epsilon X)$ and $\widetilde{\partial A}(\widetilde{\chi}) = \partial A(\chi) = \partial A(\widetilde{\chi} - \epsilon X)$ [7]. Equation (82) should be transformed into a partial differential equation before we apply it in deriving conservation laws. The symmetry condition is

$$\left[\frac{d\mathcal{L}}{d\epsilon}\right]_{\epsilon=0} \equiv 0, \tag{83}$$

which is equivalent to

$$\begin{aligned} 0 &= \int d\tau \left[\frac{d\left(\chi^\mu + \epsilon X^\mu\right)}{d\epsilon} \frac{\partial \hat{\mathcal{L}_P}}{\partial\left(\chi^\mu + \epsilon X^\mu\right)} + \sum_{s,p} \frac{d\left(\chi^\mu_{sp} + \epsilon X^\mu\right)}{d\epsilon} \frac{\partial \hat{\mathcal{L}_P}}{\partial\left(\chi^\mu_{sp} + \epsilon X^\mu\right)}\right]_{\epsilon=0} \\ &= X^\mu \int \left(\frac{\partial \hat{\mathcal{L}_P}}{\partial \chi^\mu} + \sum_{s,p} \frac{\partial \hat{\mathcal{L}_P}}{\partial \chi^\mu_{sp}}\right) d\tau. \end{aligned} \tag{84}$$

Therefore,

$$\int \left[\frac{\partial \hat{\mathcal{L}_P}}{\partial \chi^\mu} + \sum_{s,p} \frac{\partial \hat{\mathcal{L}_P}}{\partial \chi^\mu_{sp}}\right] d\tau = 0, \tag{85}$$

or

$$\frac{\partial \mathcal{L}}{\partial \chi^{\mu}}+\sum_{s,p}\int \frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^{\mu}_{sp}}d\tau=0 \tag{86}$$

Due to the integral, Eq. (86) indicates that the corresponding vector field of this symmetry for the particle-field systems is infinite dimensional.

Next, we establish the connection between the symmetry of the particle-field system, i.e., Eq. (86), and geometric conservation laws. The first term in Eq. (86) is

$$\frac{\partial \mathcal{L}}{\partial \chi^{\mu}}=\frac{D\mathcal{L}}{D\chi^{\mu}}-\frac{DA_{\nu}}{D\chi^{\mu}}\frac{\partial \mathcal{L}}{\partial A_{\nu}}-\frac{D\left(\partial_{\sigma}A_{\nu}\right)}{D\chi^{\mu}}\frac{\partial \mathcal{L}}{\partial\left(\partial_{\sigma}A_{\nu}\right)}=\frac{D}{D\chi^{\nu}}\left[\mathcal{L}\eta_{\mu}^{\ \nu}-\partial_{\mu}A_{\sigma}\frac{\partial \mathcal{L}}{\partial\left(\partial_{\nu}A_{\sigma}\right)}\right], \tag{87}$$

where use is made of the EL equation, i.e., Eq. (67). For second term in Eq. (86), we use the geometric weak EL equation (78) to obtain

$$\begin{aligned}\int_{p_1}^{p_2}\frac{\partial \hat{\mathcal{L}}_P}{\partial \chi^{\mu}_{sp}}d\tau=&\frac{D}{D\chi^{\nu}}\left\{\int\left[\left(m_sc^2+\frac{q_s}{c}A_{\sigma}\dot{\chi}^{\sigma}\right)\eta_{\mu}^{\ \nu}\delta_2-\left(\frac{q_s}{c}A_{\mu}+m_s\dot{\chi}_{sp\mu}\right)\dot{\chi}^{\nu}_{sp}\delta_2\right]d\tau\right\}\\&+\left[\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^{\mu}_{sp}}-\frac{1}{c^2}\left(\chi^{\nu}_{sp}\frac{\partial \hat{\mathcal{L}}_P}{\partial \dot{\chi}^{\mu}_{sp}}-\hat{\mathcal{L}}_P\right)\dot{\chi}_{sp\mu}\right]_{p_1}^{p_2}.\end{aligned} \tag{88}$$

The boundaries, i.e., p_1 and p_2 of the intergal of Eq. (88) should be extended to infinity, which will make the last term vanishes because of the existence of the Dirac delta function in $\hat{\mathcal{L}}_P$.

Bringing Eq. (87) and Eq. (88) into Eq. (86), we obtain

$$\frac{D}{D\chi^{\nu}}\left[\sum_{s,p}m_s\int\dot{\chi}_{sp\mu}\dot{\chi}^{\nu}_{sp}\delta_2ds-\frac{1}{2\pi}\partial_{\mu}A_{\sigma}\partial^{[\nu}A^{\sigma]}-\frac{1}{4\pi}A_{\mu}\partial^{\sigma}F_{\sigma}^{\ \nu}+\frac{1}{16\pi}F_{\sigma\rho}F^{\sigma\rho}\eta_{\mu}^{\ \nu}\right]=0, \tag{89}$$

which is equivalent to

$$\frac{D}{D\chi^{\nu}}\left\{\sum_{s,p}m_s\int\dot{\chi}_{sp\mu}\dot{\chi}^{\nu}_{sp}\delta_2d\tau+\frac{1}{4\pi}\left(-F^{\mu\sigma}F_{\sigma}^{\ \nu}+\frac{1}{4\pi}\eta^{\mu\nu}F_{\sigma\rho}F^{\sigma\rho}\right)-\frac{1}{4\pi}\frac{D}{D\chi^{\mu}}\left[\partial_{\sigma}\left(A^{\mu}F^{\nu\sigma}\right)\right]\right\}=0, \tag{90}$$

with the identity

$$-\frac{1}{2\pi}\partial_{\mu}A_{\sigma}\partial^{[\nu}A^{\sigma]}-\frac{1}{4\pi}A_{\mu}\partial^{\sigma}F_{\sigma}^{\ \nu}=-\frac{1}{4\pi}F_{\mu\sigma}F^{\nu\sigma}-\frac{1}{4\pi}\partial_{\sigma}A_{\mu}F^{\nu\sigma}. \tag{91}$$

The last term in Eq. (90) is zero because

$$-\frac{1}{4\pi}\frac{D}{D\chi^{\nu}}\frac{D}{D\chi^{\mu}}\left[\partial_{\sigma}\left(A^{\mu}F^{\nu\sigma}\right)\right]=-\frac{1}{4\pi}\frac{D}{D\chi^{(\nu}}\frac{D}{D\chi^{\sigma)}}\left[\partial_{\mu}\left(A^{\mu}F^{[\nu\sigma]}\right)\right]\equiv 0, \tag{92}$$

where $(\nu\ \mu)$ is the total symmetrization of the indexes of ν and μ [4]. Finally, we arrive at the geometric, or manifestly covariant, energy-momentum conservation laws

$$\frac{\partial}{\partial\chi^{\nu}}\left\{\sum_{s,p}m_{s}\int\dot{\chi}_{sp\mu}\dot{\chi}_{sp}^{\nu}\delta_{2}ds+\frac{1}{4\pi}\left(-F^{\mu\sigma}F_{\ \sigma}^{\nu}+\frac{1}{4\pi}\eta^{\mu\nu}F_{\sigma\rho}F^{\sigma\rho}\right)\right\}=0, \tag{93}$$

where we regard the field as defined on χ, and $\partial/\partial\chi^{\nu}\equiv D/D\chi^{\nu}$. In terms of energy-momentum tensor, Eq. (93) is

$$\partial_{\nu}T^{\mu\nu}=0, \tag{94}$$

where

$$T^{\mu\nu}=T_{P}^{\mu\nu}+T_{F}^{\mu\nu}, \tag{95}$$

$$T_{P}^{\mu\nu}=\sum_{s,p}m_{s}\int\dot{\chi}_{sp}^{\mu}\dot{\chi}_{sp}^{\nu}\delta_{2}d\tau, \tag{96}$$

$$T_{F}^{\mu\nu}=\frac{1}{4\pi}\left(-F^{\mu\sigma}F_{\ \sigma}^{\nu}+\frac{1}{4\pi}\eta^{\mu\nu}F_{\sigma\rho}F^{\sigma\rho}\right). \tag{97}$$

Here, $T_{P}^{\mu\nu}$, $T_{F}^{\mu\nu}$ and $T^{\mu\nu}$ are the energy-momentum tensors of the particles, electromagnetic field and the particle-field systems written in the geometric form, respectively. The energy-momentum tensor $T_{F}^{\mu\nu}$ for electromagnetic field described by Eq. (97) is well-known. However, to the best of our knowledge, the geometric energy-momentum tensor for particles given in Eq. (96) has not been derived previously. Instead, the energy-momentum tensor for particle is typically written in the existing literature [6, 9, 12] as

$$T_{P}^{\mu\nu}=\sum_{s,p}m_{s}\delta\left(\mathbf{X}_{sp}(t)-\boldsymbol{x}\right)\dot{\chi}_{sp}^{\mu}(t)\dot{\chi}_{sp}^{\nu}(t)\frac{d\tau(t)}{dt}, \tag{98}$$

where a Lorentzian coordinate system $\{t,\boldsymbol{x}\}$ is chosen. Obviously, Eq. (98) is not a geometric form as Eq. (96). We can prove that Eq. (96) recovers Eq. (98) when a coordinate system is chosen. For this purpose, we first show that

$$\int\delta\left(\chi_{sp}-\chi\right)g\left(\chi_{sp}\right)ds=\gamma_{sp}^{-1}(t)\delta\left(\mathbf{X}_{sp}(t)-\boldsymbol{x}\right)g\left(\chi_{sp}\left(t\right)\right), \tag{99}$$

where $\gamma_{sp}^{-1}(t)\equiv\sqrt{1-\dot{\mathbf{x}}_{sp}^{2}\left(t\right)/c^{2}}$, and $g\left(\chi_{sp}\right)$ is an arbitrary field. The left-hand side of

Eq. (99) is

$$\begin{aligned}
&\int \delta\left(\chi_{sp}-\chi\right) g\left(\chi_{sp}\right) ds \\
&\equiv \int \delta\left[\chi_{sp}\left(s_{sp}\right)-\chi\right] g\left[\chi_{sp}\left(s_{sp}\right)\right] ds_{sp} \qquad (100) \\
&= \int \delta\left[\chi_{sp}\left(s_{sp}\left(t_{sp}\right)\right)-\chi\right] g\left[\chi_{sp}\left(s_{sp}\left(t_{sp}\right)\right)\right] \frac{ds_{sp}}{dt_{sp}} dt_{sp} \\
&= \int \delta\left[c\left(t_{sp}-t\right)\right] \delta\left[\mathbf{X}_{sp}\left(t_{sp}\right)-\boldsymbol{x}\right] g\left[\chi_{sp}\left(t_{sp}\right)\right] \frac{c d\tau_{sp}\left(t_{sp}\right)}{dt_{sp}} dt_{sp}, \qquad (101)
\end{aligned}$$

where t_{sp} is the time parameter for each world-line, and $\boldsymbol{x}_{sp}\left(t_{sp}\right)$ is the space position for sp-particle at time t_{sp}. Because

$$\frac{d\tau_{sp}\left(t_{sp}\right)}{dt_{sp}} = \gamma_{sp}^{-1}(t_{sp}), \qquad (102)$$

and

$$\delta\left[c\left(t_{sp}-t\right)\right] = \frac{1}{c}\delta\left(t_{sp}-t\right), \qquad (103)$$

we have

$$\begin{aligned}
\int \delta\left(\chi_{sp}-\chi\right) g\left(\chi_{sp}\right) ds &= \int \gamma_{sp}^{-1}(t_{sp}) \delta\left[\mathbf{X}_{sp}\left(t_{sp}\right)-\boldsymbol{x}\right] g\left[\chi_{sp}\left(t_{sp}\right)\right] \delta\left(t_{sp}-t\right) dt_{sp} \\
&= \gamma_{sp}^{-1}(t) \delta\left[\mathbf{X}_{sp}\left(t\right)-\boldsymbol{x}\right] g\left[\chi_{sp}\left(t\right)\right], \qquad (104)
\end{aligned}$$

which is Eq. (99). If we take $g\left(\chi_{sp}\right) = m_s \dot{\chi}_{sp\mu} \dot{\chi}_{sp}^{\nu}$, then the geometric energy-momentum tensor for particles is

$$\begin{aligned}
T_P^{\mu\nu} &= \sum_{s,p} m_s \int \delta\left(\chi_{sp}-\chi\right) \dot{\chi}_{sp\mu} \dot{\chi}_{sp}^{\nu} ds = m_s \gamma_{sp}^{-1}(t) \delta\left[\mathbf{X}_{sp}\left(t\right)-\boldsymbol{x}\right] \dot{\chi}_{sp\mu}\left(t\right) \dot{\chi}_{sp}^{\nu}\left(t\right) \\
&= \sum_{s,p} m_s \delta\left[\mathbf{X}_{sp}\left(t\right)-\boldsymbol{x}\right] \dot{\chi}_{sp\mu}\left(t\right) \dot{\chi}_{sp}^{\nu}\left(t\right) \frac{d\tau_{sp}\left(t\right)}{dt}. \qquad (105)
\end{aligned}$$

This confirms that the geometric energy-momentum tensor for particles recovers the non-geometric form in a chosen coordinate system.

VI. CONCLUSIONS

In this paper, we have developed a manifestly covariant, or geometric, field theory for relativistic classical particle-field system, and geometrically demonstrated the connection

between space-time symmetry and energy-momentum conservation laws. In our theoretical formalism, space and time are treated with equal footing, i.e., space-time is treated as one identity without choosing a coordinate system. This is different from existing field theories where it is necessary to split space and time coordinates at certain stage, and thus the manifestly covariant property is lost.

There are several unique features in the geometric field theory developed. The first is the mass-shell condition, which induces two new terms in the geometric EL equation (51) for particles. The geometric Lagrangian density of particle-field systems is a functional of particles' world-lines (see equation (56)), which makes the symmetry vector field of the systems lies on the infinitive dimensional space (see Eq. (88)). Another feature of the theory is that particles and fields reside on different manifolds. The domain of the particle field can be the proper time or any other parameterization for the world-lines, and the electromagnetic field, on the other hand, are defined on space-time. In order to establish geometrically the connection between symmetries and energy-momentum conservations, a geometric weak EL equation (77) for particles is derived. Combining the EL equation (67) for field and the geometric weak EL equation (77) for particles, symmetries and conservation laws could be established geometrically. Using the theory, we derived for the first time a geometric expression for the energy-momentum tensor for particles in Eq. (96), which recovers the non-geometric form in the existing literature [6, 9, 12] for a chosen coordinate system.

ACKNOWLEDGMENTS

This research is supported by National Magnetic Confinement Fusion Energy Research Project (2015GB111003, 2014GB124005), National Natural Science Foundation of China (NSFC-11575185, 11575186, 11305171), JSPS-NRF-NSFC A3 Foresight Program (NSFC-11261140328), Key Research Program of Frontier Sciences CAS (QYZDB-SSW-SYS004), and the GeoAlgorithmic Plasma Simulator (GAPS) Project.

---

[1] E. Noether, Transport Theory and Statistical Physics **1**, 186 (1971).

[2] P. J. Olver, "Application of lie groups to differential equations," (Springer-Verlag, New York, 2007) pp. 242–283.

[3] R. Penrose, "The road to reality," (Knopf, New York, 2005) pp. 357–434.

[4] R. M. Wald, "Genaral relativity," (The University of Chicago, Chicago, 1984) pp. 23–68.

[5] M. Fecko, "Differential geometry and lie groups for physicists," (Cambridge, London, 2006) pp. 93–125, 327–476.

[6] L. D. Landau and E. M. lifshitz, "The classical theory of fields," (Butterworth-Heinemann, Oxford, 1975) pp. 46–89.

[7] H. Qin, J. W. Burby, and R. C. Davidson, Phys. Rev. E **90**, 043102 (2014).

[8] L. D. Landau and E. M. Lifshitz, "Mechanics," (Butterworth-Heinemann, Oxford, 1981) pp. 1–12.

[9] A. O. Barut, "Electrodynamics and classical theory of fields & particles," (Dover Publications, New York, 1964) pp. 47–68,135–164.

[10] P. A. M. Dirac, "The principles of quantum mechanics," (Oxford University, 1978) pp. 58–61.

[11] J. E. Marsden and T. S. Ratiu, "Introduction to mechanics and symmetry," (Springer, New York, 2002) pp. 1–50.

[12] J. D. Jackson, "Classical electrodynamics," (John Wiley & Sons, New York, 1999) pp. 579–617.

[13] F. Rohrlich, "Classical charged particles," (Syracuse, New York, 2007) pp. 74–85.